

BP:	1	HR:	BPM

## **CONSENT FOR EXTRACTION OF TEETH**

Patient's Na	me:	
Please initi	al each paragraph after reading. If you have	any questions, please ask your doctor BEFORE initialing.
	e right to be informed about your diagnosis and g the risks and benefits.	planned surgery so that you can decide whether to have a procedure or not
Your diagno	sis is: $\square$ Unrestorable Tooth $\square$ Wisdom Teeth	□ Periodontitis □ Apical Abscess □ Other
Your planne	d treatment is: ☐ Extraction of teeth #(s)	
Alternative t	reatment methods include: $\square$ Root Canal Treatm	nent  None  Other
	out is a permanent process. Whether the pro They include the following and others:	cedure is easy or difficult, it is still a surgical procedure. All surgeries have
1. 2. 3. 4. 5. 6. 7.  8. 9. 10. 11. 12. 13.	Numbness, pain, or changed feelings in the to due to the closeness of tooth roots (mainly we the numbness or pain goes away, but in some Trismus – you can only open your mouth a lift happens because of jaw joint (TMJ) problems from having your mouth open wide and/or for signs. This may need separate additional treat Bleeding – oozing can often happen for several Sharp ridges or bone splinters may form later another surgery to smooth or remove. Sometimes tooth roots may be left in to avoid your upper back teeth). The roots of the upper back teeth are often copening may occur from the sinus into the moult is very rare that the jaw will break, but it is possible.	ent. Iter surgery, that may need more care. Iter surgery, that may need more care. Ites being taken out, more often those with large fillings or caps. Iteeth, gums, lip, chin and/or tongue (including possible loss of taste). This is Ith wisdom teeth) to the nerves which can be injured or damaged. Usually Ites cases, it may need more treatment or may be permanent. Ites is most common after wisdom teeth are taken out. Sometimes it It already there. Damage can occur to the ligaments of the jaw joint (TMJ) If a period of time. This is more common if you already have symptoms or It hours, but a lot of bleeding is not common. If at the edge of the hole where the tooth was taken out. These may need It harming important things such as nerves or a sinus (a hollow place above lose to the sinus and sometimes a piece of root can get into the sinus. An
INFORMAT	ION FOR FEMALE PATIENTS USING BIRTH C	ONTROL PILLS
14.	reduce the preventive effect of birth control p	ills. My doctor has told me that some antibiotics and other medications may lls, and I could conceive and become pregnant. I agree to discuss with my trol during my treatment, and to continue those methods until my personal bral birth control pills.
CONSENT		
of treatmen understand including al	t or no treatment at all are choices I have an the above and give my consent to surgery and	vill be perfect. I understand that the treatment listed above and other forms d the risks of those choices have been presented to me. I have read and chosen anesthesia. I have given a complete and truthful medical history, ify that I speak, read and write English. All of my questions have been
Patient's (or	Legal Guardian's) Signature:	Date:
Doctor's Sig	nature:	Date:
Witness' Sig	uneture	Date